

CURRICULUM VITAE

Maoto Habara M,D.

Cardiovascular medicine, Toyohashi Heart Center
21-1 Gobudori, Oyama-cho, Toyohashi 441-8530 JAPAN
Office telephone number: +81-532-37-3377
Office fax number: +81-532-37-3366
habara@heart-center.or.jp

Sex: Male

Date of Birth: August 5, 1976

Age.46

Place of Birth: Hiroshima prefecture, Japan

Nationality: Japan

➤ Professional Experience

- Graduate school of medicine, Ehime University, March 2001
- Resident at Department of Cardiovascular Medicine, Okayama University, Okayama, July 2001 to August 2001
- Resident of Internal medicine at Hiroshima City Hospital - September 2001 to August 2003
- Resident of Cardiovascular Medicine, Matsuyama City Hospital – September 2003 to March 2005
- Regular staff of Cardiovascular Medicine, Kochi health science center – April 2005 to March 2007
- Regular staff of Cardiovascular Medicine, Toyohashi Heart Center – April 2007 to March 2014
- Chief physician of Cardiovascular Medicine, Toyohashi Heart Center- April 2014 to March 2020
- Co-director of Cardiovascular Medicine, Toyohashi Heart Center- April 2020 to now

- **Professional Specialty:** Interventional cardiology
- **Professional Society.**
 - Member of The Japanese Society of Internal Medicine
 - The Japan Circulation Society
 - The Japanese Society of Interventional Cardiology
 - The Japanese College of Cardiology
 - The Japanese Society of Echocardiography
- ◆ Representative manager of Japanese Retrograde Summit
- ◆ Coronary Course Directors of CCT
- ◆ Course Directors of CTO-Club

➤ Publications

Habara M, Tsuchikane E, Shimizu K, Kashima Y, Shimoji K, Nakamura S, Niizeki T, Tsutsumi T, Ito Y, Kawasaki T; Bridge point Club Japan. Japanese multicenter registry evaluating the antegrade dissection reentry with cardiac computerized tomography for chronic coronary total occlusion. *Cardiovasc Interv Ther*. 2022 Jan;37(1):116-127

Habara M, Tsuchikane E, Nasu K, Kinoshita Y, Murata A, Suzuki Y, Kawase Y, Okubo M, Matsuo H, Suzuki T. Efficacy of plaque debulking for bifurcated or ostial lesions by directional coronary atherectomy prior to second-generation drug eluting stenting. *Catheter Cardiovasc Interv*. 2020 Oct 1;96(4):773-781.

Habara M, Otsuka F, Tsuchikane E, Terashima M, Nasu K, Kinoshita Y, Murata A, Suzuki Y, Kawase Y, Okubo M, Matsuo H, Matsubara T, Yasuda S, Ishibashi-Ueda H, Suzuki T. In vivo tissue characterization of human atherosclerotic plaques by optical coherence tomography: A directional coronary atherectomy study with histopathologic confirmation. *Int J Cardiol*. 2018 1;268:1-10.

Habara M, Tsuchikane E, Nasu K, Kinoshita Y, Yaguchi T, Suzuki Y, Matsuo H, Terashima M, Matsubara T, Suzuki T. The first clinical experience with a novel

directional coronary atherectomy catheter: Preliminary Japanese multicenter experience. *Catheter Cardiovasc Interv.* 2017;89(5):880-887

Habara M, Tsuchikane E, Muramatsu T, Kashima Y, Okamura A, Mutoh M, Yamane M, Oida A, Oikawa Y, Haegawa K, for the Retrograde Summit Investigators: Comparison of Percutaneous Coronary Intervention for Chronic Total Occlusion Outcome According to Operator Experience from the Japanese Retrograde Summit Registry. *Catheter Cardiovasc Interv.* 2016;87(6):1027-35

Habara M, Nasu K, Terashima M, Ko E, Yokota D, Ito T, Kurita T, Teramoto T, Kimura M, Kinoshita Y, Tsuchikane E, Asakura Y, Matsubara T, Suzuki T: Impact on optical coherence tomographic coronary findings of fluvastatin alone versus fluvastatin + ezetimibe. *Am J Cardiol.* 2014 Feb 15;113(4):580-7.

Habara M, Terashima M, Nasu K, Kaneda H, Yokota D, Ito T, Kurita T, Teramoto T, Kimura M, Kinoshita Y, Tsuchikane E, Asakura Y, Suzuki T. Morphological differences of tissue characteristics between early, late, and very late restenosis lesions after first generation drug-eluting stent implantation: an optical coherence tomography study. *Eur Heart J Cardiovasc Imaging.* 2013;14(3):276-84

Habara M, Nasu K, Terashima M, Kaneda H, Yokota D, Ko E, Ito T, Kurita T, Tanaka N, Kimura M, Ito T, Kinoshita Y, Tsuchikane E, Asakura K, Asakura Y, Katoh O, Suzuki T. Response to letter regarding article, "impact of frequency-domain optical coherence tomography guidance for optimal coronary stent implantation in comparison with intravascular ultrasound guidance". *Circ Cardiovasc Interv.* 2012; 1;5(4):e60.

Habara M, Nasu K, Terashima M, Kaneda H, Yokota D, Ko E, Ito T, Kurita T, Tanaka N, Kimura M, Ito T, Kinoshita Y, Tsuchikane E, Asakura K, Asakura Y, Katoh O, Suzuki T. Impact of frequency-domain optical coherence tomography guidance for optimal coronary stent implantation in comparison with intravascular ultrasound guidance. *Circ Cardiovasc Interv.* 2012 Apr;5(2):193-201.

Habara M, Terashima M, Nasu K, Kaneda H, Inoue K, Ito T, Kamikawa S, Kurita T, Tanaka N, Kimura M, Kinoshita Y, Tsuchikane E, Matsuo H, Ueno K, Katoh O, Suzuki T. Difference of tissue characteristics between early and very late restenosis lesions

after bare-metal stent implantation: an optical coherence tomography study. *Circ Cardiovasc Interv.* 2011 Jun;4(3):232-8.

Habara M, Kinoshita Y, Suzuki T. Novel use of a local drug delivery catheter for coronary perforation. *J Invasive Cardiol.* 2011 Jan;23(1): 236-9.

Habara M, Terashima M, Suzuki T.: Detection of atherosclerotic progression with rupture of degenerated in-stent intima five years after bare-metal stent implantation using optical coherence tomography. *J Invasive Cardiol.* 2009 21(10):552-3.

Habara M, Kinoshita Y.: Stenting with PRotruding strut INTo side branch (SPRINT) technique: a novel technique for coronary bifurcation stenting. *Catheter Cardiovasc Interv.* 2009;73(3):336-43.

Habara M, Fujieda H, Nakamura Y. Images in cardiology. Atrial myocarditis: a possible cause of idiopathic enlargement of bilateral atria. *Heart.* 2006 ;92(6):842

Tan M, Urasawa K, Haraguchi T, Ando H, Tsubakimoto Y, Kamoi D, **Habara M**, Nakama T, Sakamoto H, Kimura M, Doijiri T, Sugihara M, Imoto Y, Sato T, Hayashi T, Sato Y. Mortality risk after use of a paclitaxel-coated stent in femoropopliteal peripheral artery disease.

Cardiovasc Interv Ther. 2022 Jan;37(1):136-144.

Konishi H, **Habara M**, Nasu K, Koshida R, Kinoshita Y, Tsuchikane E, Terashima M, Matsubara T, Suzuk, Impact of optimal preparation before drug-coated balloon dilatation for de novo lesion in patients with coronary artery disease. *Cardiovasc Revasc Med.* 2021 Mar 20:S1553-8389(21)00146-9.

Terashima M, Kaneda H, Honda Y, Shimura T, Kodama A, **Habara M**, Suzuki T: Current status of hybrid intravascular ultrasound and optical coherence tomography catheter for coronary imaging and percutaneous coronary intervention. *J Cardiol.* 2021 May;77 (5) ;435-443

Shimizu K, Shimura T, Terashima M, Yamamoto M, Adachi Y, Kano S, **Habara M**, Nasu K, Koshida R, Kinoshita Y, Tsuchikane E, Matsubara T, Suzuki T: Incidence and

predictive factors of chronic vessel enlargement after percutaneous coronary intervention for severely stenotic and occluded lesions. *Cardiovasc Revasc Med.* 2020 Oct 24:S1553-8389(20)30664-3. Online ahead of print.

Katoh H, Yamane M, Muramatsu T, Okamura A, Kashima Y, Matsuno S, Sakurada M, Kijima M, Tanabe M, **Habara M**. Safety of percutaneous coronary intervention for chronic total occlusion in patients with multi-vessel disease; sub-analysis of the Japanese Retrograde Summit Registry. *Cardiovasc Revasc Med.* 2020 Oct 20. Online ahead of print.

Aoki J, Nakazawa G, Ando K, Nakamura S, Tobaru T, Sakurada M, Okada H, Hibi K, Zen K, Ikuta A, Fujii K, **Habara M**, Ako J, Asano T, Ozaki S, Fusazaki T, Kozuma K; ELEGANT investigators. Impact of hemodialysis on clinical and angiographic outcomes in in-stent restenotic lesions following optical coherence tomography-guided drug-coated balloon treatment. *Cardiovasc Interv Ther.* 2020 Oct 13. Online ahead of print.

Ebisawa S, Kohsaka S, Muramatsu T, Kashima Y, Okamura A, Yamane M, Sakurada M, Matsuno S, Kijima M, **Habara M**. Derivation and validation of the J-CTO extension score for pre-procedural prediction of major adverse cardiac and cerebrovascular events in patients with chronic total occlusions. *PLoS One.* 2020 Sep 11;15(9):e0238640.

Kitani S, Igarashi Y, Tsuchikane E, Nakamura S, Seino Y, **Habara M**, Takeda Y, Shimoji K, Yasaka Y, Kijima M. Efficacy of drug-coated balloon angioplasty after directional coronary atherectomy for coronary bifurcation lesions (DCA/DCB registry). *Catheter Cardiovasc Interv.* 2020 Aug 10. Online ahead of print.

Nakachi T, Kohsaka S, Yamane M, Muramatsu T, Okamura A, Kashima Y, Matsuno S, Sakurada M, Seino Y, **Habara M**. Scoring System for Identification of "Survival Advantage" after Successful Percutaneous Coronary Intervention in Patients with Chronic Total Occlusion. *J Clin Med.* 2020 May 2;9(5):1319.

Aoki J, Nakazawa G, Ando K, Nakamura S, Tobaru T, Sakurada M, Okada H, Hibi K, Zen K, Habara S, Fujii K, **Habara M**, Ako J, Asano T, Ozaki S, Fusazaki T, Kozuma K; ELEGANT Investigators. Effect of combination of non-slip element balloon and drug-coating balloon for in-stent restenosis lesions (ELEGANT study). *J Cardiol.* 2019 Jun 24. pii: S0914-5087(19)30138-8.

Lee PH, Lee SW, **Habara M**, Nasu K. Unexpected Stenosis within Significantly Enlarged Distal Vessel after Successful Coronary Chronic Total Occlusion Recanalization. *Korean Circ J.* 2019 Apr;49(4):366-368.

Takeuchi W, **Habara M**, Tsuchikane E, Suzuki T. The antegrade dissection and re-entry technique as preparation of intravascular ultrasound guided re-wiring. *Cardiovasc Interv Ther.* 2019 Feb 26. [Epub ahead of print]

Yamamoto M, Tsuchikane E, Kagase A, Shimura T, Teramoto T, Kimura M, Matsuo H, Kawase Y, Suzuki Y, Kano S, **Habara M**, Nasu K, Kinoshita Y, Terashima M, Matsubara T, Suzuki T. Novel proctorship effectively teaches interventionists coronary artery chronic total occlusion lesions. *Cardiovasc Revasc Med.* 2017 Oct 26.

Nakachi T, Kohsaka S, Yamane M, Muramatsu T, Okamura A, Kashima Y, Matsuno S, Sakurada M, Kijima M, Tanabe M, **Habara M**. Impact of Hemodialysis on Procedural Outcomes of Percutaneous Coronary Intervention for Chronic Total Occlusion: Insights From the Japanese Multicenter Registry. *J Am Heart Assoc.* 2017 Oct 11;6(10).

Nasu K, Oikawa Y, **Habara M**, Shirai S, Abe H, Kadotani M, Gotoh R, Hozawa H, Ota H, Suzuki T, Shibata Y, Tanabe M, Nakagawa Y, Serikawa T, Nagasaka S, Takeuchi Y, Fujimoto Y, Tamura H, Kobori Y, Yajima J, Aizawa T, Suzuki T; NO-RECOIL Registries Investigators. Efficacy of biolimus A9-eluting stent for treatment of right coronary ostial lesion with intravascular ultrasound guidance: a multi-center registry. *Cardiovasc Interv Ther.* 2017. Epub ahead of print

Teramoto T, Tsuchikane E, Yamamoto M, Matsuo H, Kawase Y, Suzuki Y, Kanou S, Shimura T, Sato H, **Habara M**, Nasu K, Kimura M, Kinoshita Y, Terashima M, Matsubara T, Suzuki T. Successful revascularization improves long-term clinical outcome in patients with chronic coronary total occlusion. *Int J Cardiol Heart Vasc.* 2016;14:28-32

Shimura T, Yamamoto M, Tsuchikane E, Teramoto T, Kimura M, Matsuo H, Kawase Y, Suzuki Y, Kano S, **Habara M**, Nasu K, Kinoshita Y, Terashima M, Matsubara T, Suzuki T. Rates of future hemodialysis risk and beneficial outcomes for patients with

chronic kidney disease undergoing recanalization of chronic total occlusion. *Int J Cardiol.* 2016 222:707-13

Shimura T, Yamamoto M, Tsuchikane E, Teramoto T, Kimura M, Satou H, Matsuo H, Kawase Y, Suzuki Y, Kano S, **Habara M**, Nasu K, Kinoshita Y, Terashima M, Matsubara T, Suzuki T. Safety of Live Case Demonstrations in Patients Undergoing Percutaneous Coronary Intervention for Chronic Total Occlusion. *Am J Cardiol.* 2016 Oct 1;118(7):967-7

Ito T, Tsuchikane E, Nasu K, Suzuki Y, Kimura M, Ehara M, Terashima M, Kinoshita Y, **Habara M**, Suzuki T, Ohte N. Impact of lesion morphology on angiographic and clinical outcomes in patients with chronic total occlusion after recanalization with drug-eluting stents: a multislice computed tomography study. *Eur Radiol.* 2015 Oct;25(10):3084-92

Ishizuka S, **Habara M**, Nasu K. Kissing iliac artery stent technique for salvage of a total occlusion of a jailed common iliac artery. *J Invasive Cardiol.* 2014 Mar;26(3):E29-31.

Ito T, Kimura M, Ehara M, Terashima M, Nasu K, Kinoshita Y, **Habara M**, Tsuchikane E, Suzuki T. Impact of sirolimus-eluting stent fractures without early cardiac events on long-term clinical outcomes: a multislice computed tomography study. *Eur Radiol.* 2014 May;24(5):1006-12.

Teramoto T, Tsuchikane E, Matsuo H, Suzuki Y, Ito T, Ito T, **Habara M**, Nasu K, Kimura M, Kinoshita Y, Terashima M, Asakura Y, Matsubara T, Suzuki T. Initial success rate of percutaneous coronary intervention for chronic total occlusion in a native coronary artery is decreased in patients who underwent previous coronary artery bypass graft surgery. *JACC Cardiovasc Interv.* 2014 Jan;7(1):39-46.

Kubo T, Akasaka T, Shite J, Suzuki T, Uemura S, Yu B, Kozuma K, Kitabata H, Shinke T, **Habara M**, Saito Y, Hou J, Suzuki N, Zhang S: OCT compared with IVUS in a coronary lesion assessment: the OPUS-CLASS study. *JACC Cardiovasc Imaging.* 2013 ;6(10):1095-104.

Nasu K, Terashima M, **Habara M**, Ko E, Ito T, Yokota D, Ishizuka S, Kurita T, Kimura M, Kinoshita Y, Asakura Y, Tsuchikane E, Katoh O, Suzuki T. Impact of cholesterol metabolism on coronary plaque vulnerability of target vessels: a combined analysis of virtual histology intravascular ultrasound and optical coherence tomography. *JACC Cardiovasc Interv.* 2013;6(7):746-55.

Tsuchikane E, Kimura M, Suzuki T, **Habara M**, Kurita T, Tanaka N, Nasu K, Ito T, Kinoshita Y, Wyman RM. New re-entry device for revascularization of chronic coronary total occlusions: preliminary single Japanese center experience. *J Invasive Cardiol.* 2012;24(8):396-400.

Ko E, Nasu K, **Habara M**, Tanaka N, Kimura M, Kinoshita Y, Tsuchikane E, Asakura Y, Katoh O, Suzuki T. Long-term serial angiographic outcomes after sirolimus-eluting stent implantation. *Catheter Cardiovasc Interv.* 2013; 1;81(1):E29-35

Terashima M, Kaneda H, Nasu K, Matsuo H, **Habara M**, Ito T, Tanaka N, Rathore S, Kinoshita Y, Kimura M, Ehara M, Suzuki Y, Suzuki T. Protective effect of telmisartan against endothelial dysfunction after coronary drug-eluting stent implantation in hypertensive patients. *JACC Cardiovasc Interv.* 2012;5(2):182-90.

Ito T, Terashima M, Kaneda H, Nasu K, Ehara M, Kinoshita Y, Ito T, Kimura M, Tanaka N, **Habara M**, Tsuchikane E, Suzuki T. In vivo assessment of ergonovine-induced coronary artery spasm by 64-slice multislice computed tomography. *Circ Cardiovasc Imaging.* 2012;5(2):226-32.

Ito T, Matsuo H, Terashima M, Nasu K, Kinoshita Y, Ehara M, **Habara M**, Tsuchikane E, Suzuki T. Turbulent flow inside coronary artery mimicking acute coronary syndrome in multislice computed tomography. *Int J Cardiol.* 2012;31;157(2):e38-9.

Ito T, Terashima M, Kaneda H, Nasu K, Matsuo H, Ehara M, Kinoshita Y, Kimura M, Tanaka N, **Habara M**, Katoh O, Suzuki T. Comparison of in vivo assessment of vulnerable plaque by 64-slice multislice computed tomography versus optical coherence tomography. *Am J Cardiol.* 2011; 1;107(9):1270-7.

Ito T, Terashima M, Kaneda H, Nasu K, Kinoshita Y, **Habara M**, Tsuchikane E, Suzuki T. Very late sirolimus-eluting stent thrombosis due to stent fracture and late-acquired

incomplete stent apposition detected on multislice computed tomography. *Circ J.* 2011;75(11):2716-7.

Rathore S, Terashima M, **Habara M**, Kinoshita Y, Nasu K, Katoh O, Suzuki T. *J Interv Cardiol.* :Incomplete stent apposition after coronary stent implantation: myth or reality? *J Interv Cardiol.* 2009;22(4):341-9.

Ehara M, Terashima M, Kawai M, Matsushita S, Tsuchikane E, Kinoshita Y, Kimura M, Nasu K, Tanaka N, Fujita H, **Habara M**, Ito T, Rathore S, Katoh O, Suzuki T.:Impact of multislice computed tomography to estimate difficulty in wire crossing in percutaneous coronary intervention for chronic total occlusion. *J Invasive Cardiol.* 2009;21(11):575-82.

Terashima M, Rathore S, Suzuki Y, Nakayama Y, Kaneda H, Nasu K, **Habara M**, Katoh O, Suzuki T.: Accuracy and reproducibility of stent-strut thickness determined by optical coherence tomography.*J Invasive Cardiol.* 2009; 21(11):602-5.

Kinoshita Y, Katoh O, Matsubara T, Ehara M, Nasu K, **Habara M**, Suzuki T.: First clinical experience of "flower petal stenting": a novel technique for the treatment of coronary bifurcation lesions.*JACC Cardiovasc Interv.* 2010;3(1):58-65.

Tanaka N, Terashima M, Rathore S, Itoh T, **Habara M**, Nasu K, Kimura M, Itoh T, Kinoshita Y, Ehara M, Tsuchikane E, Asakura K, Asakura Y, Katoh O, Suzuki T.: Different patterns of vascular response between patients with or without diabetes mellitus after drug-eluting stent implantation: optical coherence tomographic analysis. *JACC Cardiovasc Interv.* 2010;3(10):1074-9.

➤ Book

Jang YS et al. Percutaneous Coronary Interventions for Chronic Total Occlusion, 2019, pp.121-126: CrossBoss and Stingray System: A Guide to Success: **Habara M**, Lee SW, Tsuchikane E.